



Reasons to go to ER: Green, Blood or Projectile Vomiting; Difficulty Breathing; No Urination in 3-4 hours; Frustration that may lead to harm your child; Fever alone is no longer medical emergency unless your child looks sick and does not respond to Tylenol.

Bright Futures Parent Handout 6 Month Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

Feeding Your Baby

- Most babies have doubled their birth weight.
- Your baby's growth will slow down.
- If you are still breastfeeding, that's great! Continue as long as you both like.
- If you are formula feeding, use an iron-fortified formula.
- You may begin to feed your baby solid food when your baby is ready.
- Some of the signs your baby is ready for solids
 - Opens mouth for the spoon.
 - Sits with support.
 - Good head and neck control.
 - Interest in foods you eat.

Starting New Foods

- Introduce new foods one at a time.
 - Iron-fortified cereal
- Good sources of iron include
 - Red meat
- Introduce fruits and vegetables after your baby eats iron-fortified cereal or pureed meats well. **(Avoid cow's milk, honey)**
 - Offer 1–2 tablespoons of solid food 2–3 times per day.
- Avoid feeding your baby too much by following the baby's signs of fullness.
 - Leaning back
 - Turning away
- Do not force your baby to eat or finish foods.
 - It may take 10–15 times of giving your baby a food to try before she will like it.
- ~~Avoid foods that can cause allergies—peanuts, tree nuts, fish, and shellfish.~~
- To prevent choking
 - Only give your baby very soft, small bites of finger foods.
 - Keep small objects and plastic bags away from your baby.

- Use saline rinse for congestion and colds. Generally need to on 10-14 days before other therapy.
- Start brushing gums with toothbrush or wash cloth twice daily.
- Fluoride if none in water.

FAMILY FUNCTIONING

How Your Family Is Doing

- Call on others for help.
- Encourage your partner to help care for your baby.
- Ask us about helpful resources if you are alone.
- Invite friends over or join a parent group.
- Choose a mature, trained, and responsible babysitter or caregiver.
- You can talk with us about your child care choices.

Healthy Teeth

- Many babies begin to cut teeth.
- Use a soft cloth or toothbrush to clean each tooth with water only as it comes in.
- Ask us about the need for fluoride.
- Do not give a bottle in bed.
- Do not prop the bottle.
- Have regular times for your baby to eat. Do not let him eat all day.

ORAL HEALTH

Your Baby's Development

- Place your baby so she is sitting up and can look around.
- Talk with your baby by copying the sounds your baby makes.
- Look at and read books together.
- Play games such as peekaboo, patty-cake, and so big.
- Offer active play with mirrors, floor gyms, and colorful toys to hold.
- If your baby is fussy, give her safe toys to hold and put in her mouth and make sure she is getting regular naps and playtimes.
- Put your baby to bed when she is sleepy but still awake.

INFANT DEVELOPMENT

Crib/Playpen

- Lower the crib mattress all the way when your baby begins to stand.
- Use a crib with slats close together—2 3/8 inches apart or less.
- When your baby is in the crib, make sure the drop side is up.
- Don't use loose or soft bedding.
- Use a mesh playpen with weaves less than 1/4 inches apart.

Safety (Rear facing car seat until 2 years of age)

- Use a rear-facing car safety seat in the back seat in all vehicles, even for very short trips.
- Never put your baby in the front seat of a vehicle with a passenger air bag.
- Don't leave your baby alone in the tub or high places such as changing tables, beds, or sofas.
- While in the kitchen, keep your baby in a high chair or playpen.
- Do not use a baby walker.
- Place gates on stairs.
- Close doors to rooms where your baby could be hurt, like the bathroom.
- Prevent burns by setting your hot water heater so the temperature at the faucet is 120°F or lower.
- Turn pot handles inward on the stove.
- Do not leave hot irons or hair care products plugged in.
- Never leave your baby alone near water or in bathwater, even in a bath seat or ring.
 - Always be close enough to touch your baby.
- Lock up poisons, medicines, and cleaning supplies; call Poison Help if your baby eats them.

SAFETY

What to Expect at Your Baby's 9 Month Visit

We will talk about

- Disciplining your baby
- Introducing new foods and establishing a routine
- Helping your baby learn
- Car seat safety
- Safety at home

NEXT WELL- 9 Months

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org
NO TV until 2 years of age!!

NUTRITION AND FEEDING

Appointment Line and After Hours Provider:
410-293-2273

Sign up for Relay Health to e-mail
communicate with your provider at
www.RELAYHEALTH.com

Review your labs on TRICAREONLINE.COM



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.

Starting Solid Foods



Until now, your baby's diet has been made up of breast milk and/or formula. But once your baby reaches 4 to 6 months of age, you can begin adding solid foods. Read on to learn more about introducing solid foods.

When can my baby eat solid foods?

Most babies are ready to eat solid foods at 4 to 6 months of age. Before this age instead of swallowing the food, they push their tongues against the spoon or the food. This tongue-pushing reflex is necessary when they are breastfeeding or drinking from a bottle. Most babies stop doing this at about 4 months of age. Energy needs of babies begin to increase around this age as well, making this a good time to introduce solids.

You may start solid foods with any feeding. Try scheduling feedings during family meals. Or if your baby is easily distracted, you may want to pick a quiet time when you can focus on feeding your baby. However, keep in mind that as your child gets older, she will want to eat with the rest of the family.

Feeding your baby solid foods

To prevent choking, make sure your baby is sitting up when you introduce solid foods. If your baby cries or turns away when you give him the food, do not force the issue. It is more important that you both enjoy mealtimes than for your baby to start solids by a specific date. Go back to nursing or bottle-feeding exclusively for a time before trying again. Remember that starting solid foods is a gradual process and at first your baby will still be getting most of his nutrition from breast milk and/or formula.

It is important for your baby to get used to the process of eating—sitting up, taking bites from a spoon, resting between bites, and stopping when full. Always use a spoon to feed your baby solid foods. These early experiences will help your child learn good eating habits throughout life.

Some parents try putting baby cereal in a bottle. This is not a good idea. Feeding your baby this way can cause choking. It also may increase the amount of food your baby eats and can cause your baby to gain too much weight. However, cereal in a bottle may be recommended if your baby has reflux. Check with your child's doctor.

How to start

Start with half a spoonful or less and talk to your baby through the process ("Mmm, see how good this is!"). Your baby may not know what to do at first. She may look confused, wrinkle her nose, roll the food around her mouth, or reject it altogether. This is a normal early reaction to solid foods.

One way to make eating solids for the first time easier is to give your baby a little breast milk and/or formula first, then switch to very small half-spoonfuls of food, and finish with more breast milk and/or formula. This will prevent your baby from getting frustrated when she is very hungry.

Do not be surprised if most of the first few solid-food feedings wind up on your baby's face, hands, and bib. Increase the amount of food gradually, with just a teaspoonful or two to start. This allows your baby time to learn how to swallow solids.

What kinds of foods should my baby eat?

For most babies it does not matter what the first solid foods are. By tradition, single-grain cereals are usually introduced first. However, there is no medical evidence that introducing solid foods in any particular order has an advantage for your baby. Though many pediatricians will recommend starting vegetables before fruits, there is no evidence that your baby will develop a dislike for vegetables if fruit is given first. Babies are born with a preference for sweets, and the order of introducing foods does not change this. If your baby has been mostly breastfeeding, he may benefit from baby meat, which contains more easily absorbed sources of iron and zinc that are needed by 4 to 6 months of age. Please discuss this with your child's doctor.

Baby cereals are available premixed in individual containers or dry, to which you can add breast milk, formula, or water. Premixed baby cereals are convenient, while dry cereals are richer in iron and allow you to control the thickness of the cereal. Whichever type of cereal you use, make sure that it is made for babies because these cereals contain extra nutrients your baby needs at this age.

Using a high chair

The following are safety tips when using a high chair:

- Make sure the high chair you use cannot be tipped over easily.
- If the chair folds, be sure it is locked each time you set it up.
- Whenever your child sits in the chair, use the safety straps, including the crotch strap. This will prevent your child from slipping down, which could cause serious injury or even death. Never allow your child to stand in the high chair.
- Do not place the high chair near a counter or table. Your child may be able to push hard enough against these surfaces to tip the chair over.
- Never leave a young child alone in a high chair and do not allow older children to climb or play on it because this could also tip it over.
- A high chair that hooks on to a table is not a good substitute for a freestanding one. If you plan to use this type of chair when you eat out or when you travel, look for one that locks on to the table. Be sure the table is heavy enough to support your child's weight without tipping. Also, check to see whether your child's feet can touch a table support. If your child pushes against the table, it may dislodge the seat.

Once your baby learns to eat one food, gradually give him other foods. Generally, meats and vegetables contain more nutrients per serving than fruits or cereals.

Many pediatricians recommend against giving eggs and fish in the first year of life because of allergic reactions, but there is no evidence that introducing these nutrient-dense foods after 4 to 6 months of age determines whether your baby will be allergic to them. Give your baby one new food at a time, and wait at least 2 to 3 days before starting another. After each new food, watch for any allergic reactions such as diarrhea, rash, or vomiting. If any of these occur, stop using the new food and consult with your child's doctor.

Within a few months of starting solid foods, your baby's daily diet should include a variety of foods each day that may include the following:

- Breast milk and/or formula
- Meats
- Cereal
- Vegetables
- Fruits
- Eggs and fish

Finger foods

Once your baby can sit up and bring her hands or other objects to her mouth, you can give her finger foods to help her learn to feed herself. To avoid choking, make sure anything you give your child is soft, easy to swallow, and cut into small pieces. Some examples include small pieces of banana, wafer-type cookies, or crackers; scrambled eggs; well-cooked pasta; well-cooked chicken finely chopped; and well-cooked and cut up yellow squash, peas, and potatoes. Do not give your baby any food that requires chewing at this age. (See "Choking hazards.")

At each of your child's daily meals, she should be eating about 4 ounces, or the amount in one small jar of strained baby food. Limit giving your child foods that are made for adults. These foods often contain more salt and other preservatives.

If you want to give your baby fresh food, use a blender or food processor, or just mash softer foods with a fork. All fresh foods should be cooked with no added salt or seasoning. Though you can feed your baby raw bananas (mashed), most other fruits and vegetables should be cooked until they are soft. Refrigerate any food you do not use, and look for any signs of spoilage before giving it to your baby. Fresh foods are not bacteria-free, so they will spoil more quickly than food from a can or jar.

Warning: Do not feed your baby home-prepared beets, turnips, carrots, spinach, or collard greens in the first year of life.

In some parts of the country, these vegetables have large amounts of nitrates, chemicals that can cause an unusual type of anemia (low blood count) in young babies. Baby food companies are aware of this problem and screen the produce they buy for nitrates. They also avoid buying these vegetables in parts of the country where nitrates have been found. Thus it is safer to use commercially prepared forms of these foods during the first year of life.

What changes can I expect after my baby starts solids?

When your child starts eating solid foods, his stools will become more solid and variable in color. Due to the added sugars and fats, they will have a much stronger odor too. Peas and other green vegetables may turn the stool a deep-green color; beets may make it red. (Beets sometimes make urine red as well.) If your baby's meals are not strained, his stools may contain undigested pieces of food, especially hulls of peas or corn, and the skin of tomatoes or other vegetables. All of this is normal. Your child's digestive system is still immature and needs time before it can fully process these new foods. If the stools are extremely loose, watery, or full of mucus, however, it may mean the digestive tract is irritated. In this case, reduce the amount of solids and introduce them more slowly. If the stools continue to be loose, watery, or full of mucus, consult your child's doctor to see if your child has a digestive problem.

Should I give my baby juice?

Babies do not need juice. Babies younger than 6 months should not be given juice. However, if you choose to give your baby juice, do so only after 6 months of age and offer it only in a cup, not in a bottle. To help prevent tooth decay, do not put your child to bed with a bottle. If you do, make sure it contains only water.

Limit juice intake to no more than 4 ounces a day and offer it only with a meal or snack. Any more than this will reduce her appetite for other, more nutritious foods, including breast milk and/or formula. Too much juice also can cause diaper rash, diarrhea, or excessive weight gain.

Give your child extra water if she seems to be thirsty between feedings. During the hot months when your child is losing fluid through sweat, offer water 2 or more times a day. If you live in an area where the water is fluoridated, these feedings also will help prevent future tooth decay.

Good eating habits start early

Babies and small children do not know what foods they need to eat. Your job as a parent is to offer a good variety of healthy foods that are rich in the nutrients that they need. Watch your child for cues that she has had enough to eat. Do not overfeed!

Choking hazards

Do not feed children younger than 4 years round, firm foods unless they are chopped completely. Round, firm foods are common choking dangers. When infants and young children do not grind or chew their food well, they may try to swallow it whole. The following foods can be choking hazards:

- Hot dogs (including meat sticks [baby food "hot dogs"])
- Nuts and seeds
- Chunks of meat or cheese
- Whole grapes
- Popcorn
- Chunks of peanut butter
- Raw vegetables
- Fruit chunks, such as apple chunks
- Hard, gooey, or sticky candy
- Chewing gum

Begin to build good eating habits. Usually eating 5 to 6 times a day (3 meals and 2 to 3 snacks) is a good way to meet toddlers' energy needs. Children who "graze," or eat constantly, may never really feel hungry. They can have problems from eating too much or too little.

If you are concerned that your baby is overweight or becoming overweight, talk with your child's doctor before making any changes to his diet. During these months of rapid growth, your baby needs a balanced diet that includes fat, carbohydrates, and protein. Continue to give breast milk and/or formula for the first year. After 1 year of age, if you have a family history of obesity, cardiovascular disease, or high cholesterol, your child's doctor may suggest using reduced fat milk. After 1 year of age you may also reduce the amount of food your child eats at each meal. However, it is important that he continue to get the balanced diet he needs. Talk with your child's doctor about this. Your child's doctor will help you determine if your child is eating too much, not eating enough, or eating too much of the wrong kinds of foods.

Because prepared baby foods have no added salt, they are not a source of added salt. However, as your baby eats more and more "table foods," he will imitate the way you eat, including using salt and nibbling on salty snacks. For your child's sake as well as your own, eat a healthy diet yourself and decrease your intake of fat and salty snack foods. Provide a good role model by eating a variety of healthy, nutrient-rich foods.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

From your doctor

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
Web site—www.aap.org

Copyright © 2008
American Academy of Pediatrics
All rights reserved.

Acetaminophen (Tylenol, Store Brand) Dosing Information

****Give every 4-6 hours, as needed, no more than 5 times in 24 hours****

Weight of Child	Infant Drops Old Concentration 80mg/0.8ml	Infant Oral Suspension: New Concentration 160mg/5ml	Children's Elixir 160mg/5ml	Children's Tablets 80mg =1 tablet	Junior Strength 160 mg = 1 tablet
6-11 lbs (2.7-5 kg)	0.4 ml	1 ml	1 ml		
12-17 lbs (5.5-7.7 kg)	0.8 ml	2.5ml	2.5 ml	1 tablet	
18-23 lbs (8.2-10.5 kg)	1.2 ml	3.75ml	3.75 ml	1 ½ tablets	
24-35 lbs (10.9-15.9 kg)	1.6 ml (2 droppers)	5 ml	5 ml	2 tablets	
36-47 lbs (16.4-21.4 kg)	2.4 ml (2.5 droppers)		7.5 ml	3 tablets	
48-59 lbs (21.8-26.8 kg)			10 ml	4 tablets	2 tablets
60-71 lbs (27.3-32.3 kg)			12.5 ml	5 tablets	2 ½ tablets
72-95 lbs (32.7-43.2 kg)			15 ml	6 tablets	3 tablets

Ibuprofen (Motrin, Advil, Store Brand) Dosing Information

****Give every 8 hours, as needed, no more than 4 times in 24 hours****

****NOT ADVISED FOR PATIENTS UNDER 6 MONTHS OF AGE****


Weight of Child	Infant Drops 50mg/1.25ml	Children's Elixir 100mg/5ml	Children's Tablets 50mg/1 tablet	Junior Strength 100mg/1 tablet
Under 11 lbs (5kg)	*NOT ADVISED			
12-17lbs (5.5-7.7kg)	1.25 ml	2.5 ml		
18-23lbs (8.2-10.5kg)	1.875 ml	3.75 ml	1 tablet	
24-35 lbs (10.9-15.9 kg)	2.5 ml	5 ml	2 tablets	
36-47 lbs (16.4-21.4 kg)	-	7.5 ml	3 tablets	
48-59 lbs (21.8-26.8 kg)	-	10 ml	4 tablets	2 tablets
60-71 lbs (27.3-32.3 kg)	-	12.5 ml	5 tablets	2 ½ tablets
72-95 lbs (32.7-43.2 kg)	-	15 ml	6 tablets	3 tablets

Diphenhydramine (Benadryl, Generic, Store Brand) Dosing Information

****Give every 6 hours as needed, no more than 4 times in 24 hours****

Weight of Child	Liquid 12.5mg/5ml	Chewable 12.5mg/tablet	Capsule 25mg/capsule
13.2-15.3 lbs	3 ml	-	-
15.4-17.5 lbs	3.5 ml	-	-
17.6-19 lbs	4 ml	-	-
20-24 lbs	3.75 ml	-	-
25-37 lbs	5 ml	1 tablet	-
38-49 lbs	7.5 ml	1 ½ tablets	-
50-99 lbs	10 ml	2 tablets	1 capsule
100+ lbs	-	4 tablets	2 capsules

2014 Recommended Immunizations for Children from Birth Through 6 Years Old



Is your family growing? To protect your new baby and yourself against whooping cough, get a Tdap vaccine in the third trimester of each pregnancy. Talk to your doctor for more details.

Shaded boxes indicate the vaccine can be given during shown age range.

NOTE: If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.

FOOTNOTES: * Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.
 § Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.

SEE BACK PAGE FOR MORE INFORMATION ON VACCINE-PREVENTABLE DISEASES AND THE VACCINES THAT PREVENT THEM.

For more information, call toll free **1-800-CDC-INFO** (1-800-232-4636) or visit <http://www.cdc.gov/vaccines>



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™