

18.0 MEDICATIONS

Last Reviewed: Dec 15

Last Revised: Dec 15

Note: Any medication not listed in this section is not approved for aviation. Contact NAMI Code 53HN if further guidance is needed.

18.1 NATOPS ON MEDICATIONS

General NATOPS (OPNAVINST 3710.7 series, chapter 8) includes the following statements on medications (Drugs):

Taking drugs prescribed by competent medical authority shall be considered sufficient cause for recommendation of grounding unless their use is specifically approved by a Flight Surgeon (or Aeromedical Examiner or Aeromedical Physician Assistant), or a waiver for specific drug use has been granted by CHNAVPERS or the Commandant of the Marine Corps. Consideration shall be given to the removal of ground support personnel from critical duties, for the duration of the drug effects, if appropriate. Medications such as antihistamines, antibiotics, tranquilizers, sleeping pills, etc., shall be discarded if all are not used during the period of medication.

Because of the possibility of adverse side effects and unpredictable reactions, the use of over-the-counter drugs by flight personnel is prohibited unless specifically approved by a Flight Surgeon (or Aeromedical Examiner or Aeromedical Physician Assistant). Ground support personnel shall be briefed on the hazards of self-medication and should be discouraged from using such drugs.

In general, all medications require temporary grounding unless specifically described here as NCD for flight duties.

18.2 ANTIMICROBIAL

Last Reviewed: Dec 15

Last Revised: Dec 15

All antibiotics *other than the following very specific exceptions* require grounding (CD). The listed exceptions do not forgive you from doing something obviously inadvisable such as allowing a sick person to fly.

Aviation personnel on the following approved antibiotics may be considered for an up chit prior to the completion of the course of therapy as long as the condition being treated has resolved in all significant aspects with no adverse reaction that might compromise safety of flight or mission completion.

ANTI-BACTERIAL MEDICATIONS:

ANTI-MALARIALS:

Refer to Aeromedical Reference and Waiver Guide (ARWG) section 17 on [Malaria](#).

FLUOROQUINOLONES:

CIPROFLOXACIN: NCD.

ISONIAZID: No waiver needed when used for TB prophylaxis as long as the member remains under close evaluation by flight surgeon. This medication causes occasional liver damage, especially above age 35. All personnel are to be monitored in accordance with current preventive and occupational medicine guidelines.

MACROLIDES:

ERYTHROMYCIN: NCD- including long-term, low-dose use for acne.

NITROFURANTOIN: CD. Waiver considered if under close observation of flight surgeon. Adverse effects include pneumonitis or peripheral neuropathy.

PENICILLINS:

AMPICILLIN, AMOXICILLIN, PENICILLIN VK, AUGMENTIN, DICLOXACILLIN: NCD.

SULFONAMIDES:

BACTRIM/SEPTRA: CD. Waivers will be considered for long term use.

TETRACYCLINES:

TETRACYCLINE, DOXYCYCLINE: NCD. (Including long-term use for acne).

MINOCYCLINE: CD. Prohibited due to possible vestibular side effects.

ANTI- FUNGAL MEDICATIONS:

GRISEOFULVIN: CD. Waivers are considered if under close observation by local flight surgeon. Watch for bone marrow suppression.

ITRACONAZOLE (SPORANOX): NCD. While not approved for chronic use, itraconazole has a safer profile than ketoconazole, and need not be used on a chronic basis to be effective. Recommended use in aviation personnel is to administer in week-long pulses each month for four to six cycles. Aviators should be grounded for the first 48 hours of each cycle.

Since it is not administered chronically, such as griseofulvin, a waiver is not required. The recommended initial treatment is over a weekend to allow return to flight duties the following Monday, thus minimizing flight schedule loss.

TERBINAFINE (LAMISIL): NCD. **Requires a 72-hour grounding period.** Terbinafine has a safer profile than ketoconazole and has a lower relapse rate than itraconazole. The recommended use in aviation personnel is to administer daily for twelve weeks. Aviators should be grounded for the first 72 hours and a waiver is not required when no side-effects exist and appropriate monitoring is performed. The recommended initial treatment is over a weekend to allow return to flight duties the following Monday, thus minimizing flight schedule loss.

ANTI-VIRAL MEDICATIONS:

ACYCLOVIR, VALACYCLOVIR: CD for continuous therapy. Waivers are considered for suppressive/prophylactic therapy. Intermittent therapy does not require a waiver. The patient should be grounded and monitored for side effects for a minimum of 3 days during the initial treatment or upon initiation of suppressive therapy. The need for suppressive therapy should be reassessed on an annual basis. Topical [acyclovir](#) is NCD.

OSELTAMIVIR (TAMIFLU), ZANAMIVIR (RELENZA)- NCD, **Requires a 72-hour grounding period.** These medications are indicated for prophylaxis and treatment of influenza A and B viruses. They can decrease the severity, duration and complications of influenza illnesses. These medications require a 72-hour grounding period following initiation of treatment to assess for adverse side effects. In the absence of flu symptoms and adverse side effects from the medications, flight duties may resume following the 72 hour grounding period. Reducing the initial grounding period to 48 hours may be considered for operational requirements with NAMI consultation.

TRUVADA® (EMTRICITABINE + TENOFOVIR) PRE-EXPOSURE PROPHYLAXIS: CD, **waivers considered for specific use on a case-by-case basis. See entry in Miscellaneous section for details.**

18.3 ANTI-HYPERLIPIDEMICS

EZETIMIBE (ZETIA): NCD. A waiver is not required. An initial grounding period for 72 hours is required to assess for idiosyncratic reactions. If used in combination with HMG-CoA reductase inhibitors, refer to the waiver guide section on Hyperlipidemia for additional guidance.

FIBRIC ACIDS:

FENOFIBRATE (TRICOR); GEMFIBROZIL (LOPID): CD. Fenofibrate (Tricor) and gemfibrozil (Lopid) are both considered disqualifying. A waiver may be considered after a 14-day ground trial of the medication without side-effects. Fenofibrate is preferred over gemfibrozil due to fewer side effects. Prior to initiating treatment, baseline lab studies must be obtained to include: lipid panel, liver function testing (ALT/AST/ALK PHOS), CBC, FBS, and CPK. These tests are to be repeated at three months, six months and then annually if the values remain stable. Evaluate for muscle aches (myalgias) at follow-up exams and measure CPK levels if clinically indicated. If fibric acid is used in combination with an HMG-CoA reductase inhibitor, refer to the waiver guide section on Hyperlipidemia for further guidance.

NIACIN: CD. No waiver.

RESINS:

CHOLESTYRAMINE: NCD if tolerated without side effects.

STATINS:

PRAVASTATIN, SIMVASTATIN, LOVASTATIN, ATORVASTATIN: NCD. HMG Co-A reductase inhibitors (pravastatin, simvastatin, lovastatin, atorvastatin, etc.) are all NCD and a waiver is not required. Refer to ARWG section on hypercholesterolemia for additional guidance. Lipid panel, liver function tests (ALT/AST/ALK PHOS), CBC, and CPK are recommended at baseline, 3, and 6 months, then annually. Liver enzyme elevations above three times normal are disqualifying.

18.4 ANTI-HYPERTENSIVES

Last Reviewed: Dec 15

Last Revised: Oct 15

ACE INHIBITORS (ACE-I):

CD. The entire family ([captopril](#), [enalapril](#), [lisinopril](#), etc.) is CD, but waiverable. The member must be grounded upon initiation of treatment. Waiver will be considered after 30 days of treatment if member's hypertension is controlled on a stable dosage of medication without evidence of side effects. If local pharmacy policy requires changing from one ACE-I to another, advise Code 53HN of the change. Refer to ARWG section on [hypertension](#) for additional guidance.

ANGIOTENSIN RECEPTOR BLOCKERS (ARB):

CD. These agents may be used if member does not tolerate an ACE-I or has some other specific medical indication for its use. The same guidelines used for ACE-I apply.

ANTIADRENERGIC AGENTS:

[DOXAZOSIN](#), [PRAZOSIN](#), others in class: CD. No Waiver. Call NAMI Code 53HN for further guidance.

BETA BLOCKERS (for hypertension only):

CD. Beta blockers are not considered for waivers for Service Groups I or II personnel. Senior officers (LCDR and above) may be waived to Service Group 3 or Class II flying duties in non-tactical aircraft. All SG 1, SG 2, or tactical NFOs are considered NPQ, no waiver recommended. Designated Naval Aircrew will be considered for a waiver. Aviation personnel on beta blockers should not pull more than 2.5 Gs, so requests for waivers should state "transport/maritime/helo aircraft only." Air traffic controllers are usually waived. When beta blockers are used, preference shall be given to cardioselective agents such as [atenolol](#).

CALCIUM CHANNEL BLOCKERS:

[AMLODIPINE \(NORVASC\)](#): CD. A second generation calcium channel blocker, [AMLODIPINE](#) may be considered for waiver for use in the control of hypertension only after failure to control the condition on other approved agents. These cases must be reviewed individually by NAMI prior to issuance of an Aeromedical Clearance notice. Local Board is not authorized to issue a clearance notice for [AMLODIPINE](#) use.

[NIFEDIPINE \(PROCARDIA\)](#): CD. No Waiver.

COMBINATION AGENTS:

CD. Combination agents may be used if the individual agents themselves are recommended for waiver. Follow the restrictions and guidelines outlined for each individual agent.

THIAZIDE DIURETICS:

[HYDROCHLOROTHIAZIDE \(for Hypertension\)](#): CD. [HYDROCHLOROTHIAZIDE](#) (HCTZ), with or without [triamterene](#) or [potassium](#) replacement, can be used as a first line agent for treatment of hypertension in designated personnel. ACE inhibitors are preferred as they have a low incidence of aeromedically significant side effects and are generally well tolerated. See [hypertension](#) section of ARWG for waiver criteria and further guidance.

18.5 IMMUNIZATIONS

GROUNDING FOR VACCINATIONS:

OPNAVINST 3710.7 series requires a 12 hour grounding period following immunizations unless otherwise specified in this document. The specific guidelines and grounding periods for each vaccination are described below. As per MANMED Article 15-77, the administration of routine immunizations that require a temporary grounding, do not require issuance of an Aeromedical Grounding Notice. This is a “self-limited” grounding period allowed in the absence of adverse side effects.

VACCINE ADVERSE EVENT REPORTING SYSTEM (VAERS)

The Vaccine Adverse Event Reporting System (VAERS) is used to report adverse events or reactions to all vaccines. VAERS, the primary U.S. vaccine safety monitoring system, encourages reporting of any unexpected or serious event occurring after any vaccination as well as adverse events occurring in persons following close contact with a vaccine recipient. An adverse event is any clinically significant medical event that occurs following administration of a vaccine. A VAERS report should be submitted even if it is not certain that the event was caused by the vaccine. Web reporting is available at <http://vaers.hhs.gov/>.

ANTHRAX

BACKGROUND: Human anthrax vaccine was developed in England and the U.S. in the 1950s and early 1960s. The vaccine is U.S. Food and Drug Administration (FDA)-licensed and has been routinely given in the U.S. since 1970.

The vaccine has an excellent safety record. The most common side effects reported are mild discomfort (localized swelling and redness at the site of injection), joint aches, and in a few cases, nausea, loss of appetite, and headaches. There is no evidence from records at the Michigan Biologic Products Institute (which is the only U.S. producer of the vaccine) that the vaccine is associated with permanent local or systemic effects.

DOSAGE AND ADMINISTRATION: The current dose schedule for the U.S. vaccine consists of 6 shots given over an 18 month schedule and an annual booster thereafter.

Contraindications for use are sensitivity to vaccine components (formalin, aluminum hydroxide, benzethonium chloride) and/or history of clinical anthrax. Pregnant women should not receive this vaccine until after delivery. The vaccine should be stored at refrigerator temperature (not frozen).

A 12 hour grounding period is recommended for the anthrax vaccination.

CHOLERA

Sale of the only licensed cholera vaccine in the United States has been discontinued, and the CDC does not currently recommend the vaccine for travelers because of the brief and incomplete immunity it offers. In lieu of vaccination, proper hygiene and food and water precautions should be carefully emphasized.

DIPHTHERIA TETANUS (DT) AND TETANUS TOXOID

This vaccine is used to prevent bacterial elaboration of toxins resulting in muscular spasm/lockjaw, which is usually found in the setting of a contaminated wound. These vaccines are toxoids and are both known to be 95% efficacious. They are given every 10 years, however if a suspicious wound is encountered, the standard is to revaccinate if more than 5 years has elapsed since the last vaccination. The dose is 0.5 cc IM. Adverse events include frequent local reactions. Hypotonic, hyporesponsive episodes, seizures, and acute encephalopathy have been reported on rare occasions. A 12 hour grounding period is recommended for this vaccination.

HEPATITIS A

This is an inactivated virus vaccine which is given as a 1.0 cc dose IM, with a booster dose 6 to 12 months later. Protective levels of antibodies are detectable in 80 to 98% of recipients 15 days after the first dose, and in 96% after one month. Protection is expected to last 20 years. No significant adverse events have been reported, although some recipients experience local injection site soreness. Transient systemic symptoms are uncommon. In the USA, the presence of anti-HAV antibodies indicating past infection and probable immunity increases from about 10% in young children to about 75% in adults more than 50 years old. A 12 hour grounding period is recommended for this vaccination.

HEPATITIS B

This is an inactivated virus vaccine which is given as a 1.0 cc IM dose, with boosters at one and 6 months. Current CDC recommendations are to immunize everyone 18 years of age or younger and adults over 18 who are at risk. The at-risk population includes health care and public safety worker who might have contact with blood or body fluids, people who have more than one sex partner in six months, sex contacts of infected people, people who inject illegal drugs, hemodialysis patients, and household contacts of people with chronic HBV infection. Contraindications to vaccination include a history of allergic reaction to either baker's yeast or the hepatitis B vaccine. Mild soreness at the injection site is seen in approximately one out of 11 children and adolescents and one out of four adults, and mild to moderate fever is seen in up to one out of 14 children and one out of 100 adults. A 12 hour grounding period is recommended for this vaccination.

INFLUENZA

INJECTABLE INACTIVATED INFLUENZA VACCINE

This vaccine is composed of inactivated whole or disrupted influenza viruses and changed annually to reflect antigenic changes in the A and B strains of the virus that is in circulation. Immunity after the standard 0.5 cc IM dose lasts about six months, so annual administration is required, ideally before the start of flu season. The vaccine is indicated in the elderly (>65), residents of chronic care facilities, those with cardiac, pulmonary or immunosuppressive diseases such as cancer and DM, and close contacts of those at risk. All active duty Navy and Marine Corps personnel are required to have one dose of this vaccine each year. The only contraindication is a bona fide history of generalized allergic reaction to the vaccine, eggs, or egg components. Effectiveness varies with how closely vaccine strains match the strains in the community, generally about 60-85%. A mild local reaction is the most common adverse effect, although some individuals have a transient mild "viral syndrome." A 12 hour grounding period is recommended for this vaccination.

FLUMIST

All active duty Navy and Marine Corps personnel are required to have one dose of influenza vaccine (IM or intranasal spray) each year. FluMist® (Influenza Virus Vaccine Live, Intranasal), is composed of live, attenuated influenza virus (LAIV) that is administered by nasal spray. It is used for the prevention of Influenza A and B in healthy adults under age 50 who are not pregnant. The 0.5mL dose is given as a 0.25mL spray in each nostril.

The immunization is less effective in those with pre-existing nasal congestion. The dose should be repeated if the patient sneezes following administration. Immunity after the standard intranasal dose declines during the year, so annual administration is required—ideally, before the start of “flu season.” There appears to be an increase in protective antibodies over time with subsequent doses. Effectiveness varies according to how closely the strains used to make the vaccine match those in the community.

The onset of symptoms after immunization usually occurs within the first 24 hours, with most symptoms presenting by the third day. The duration of symptoms is typically 1-2 days. The most common adverse effects include:

- headache 40%
- sore throat 28%
- tiredness 26%
- myalgias 17%
- cough, nasal congestion, and rhinitis 9-45%
- Less common adverse effects include chills, abdominal pain, diarrhea, vomiting, and otitis media.

A “self-limited” grounding period of 72 hours after immunization is required to assess for symptom severity. Commanding officers may return aeronautically designated personnel to duty involving flight operations in less than 72 hours on the recommendation of a flight surgeon when necessary to meet “real world” operational commitments. The presence and severity of symptoms may require the grounding of some personnel for greater than 72 hours. To minimize operational impact, commands may elect to stagger the administration of the vaccine to their personnel. For example, a command might elect to vaccinate 50% of eligible personnel one week and the remaining personnel the following week. Another option would be to schedule immunizations immediately prior to a period when no flights are scheduled (e.g., just prior to a holiday weekend).

Additional information is available via the CDC website at, <http://www.cdc.gov/flu/professionals/vaccination/>

ANTIVIRAL MEDICATIONS:

OSELTAMIVIR (TAMIFLU), ZANAMIVIR (RELENZA)-

NCD, Requires an initial 72-hour grounding period. See Antimicrobial Section 18.2.

JAPANESE ENCEPHALITIS (March 2010)

Japanese Encephalitis (JE), a mosquito-borne arboviral infection, is the leading cause of viral encephalitis in Asia with over 50,000 sporadic and epidemic cases reported annually. Two inactivated virus vaccines are currently available, JE-Vax, licensed in 1993, and Ixiaro, licensed in 2009.

JE-VAX. JE-Vax is administered as a 1.0 mL SC dose with an effectiveness of 80-90%. Intradermal dosing at two sites is as immunogenic as a single SC dose. Three doses during a 30 day period (days 0, 7, and 30) provides the longest immunogenic protection. A booster given at one year will significantly increase antibody titers, which may then persist for several years. An abbreviated schedule of immunizations given on days 0, 7, and 14 may be used if significant time constraints exist.

JE-Vax is associated with a moderate frequency of local and mild systemic side effects. About 20% of recipients experience local redness, swelling, or tenderness, and systemic side effects (fever, headache, malaise, and rash) have been reported in about 10% of vaccine recipients. An additional pattern of adverse reactions characterized by generalized urticaria and/or angioedema, and rarely respiratory distress or collapse, has been reported. These reactions occurred after a longer interval and usually after the first or second dose. The median time to onset of symptoms after the first dose is 12 hours, and 88% of reactions occur within 3 days. The interval after the second dose is longer, with a median time of 3 days and possibly as long as two weeks. After reviewing the experiences of I-MEF personnel during the first several years of use, the original 3-5-3 day grounding regimen appears excessive based upon the actual observed incidence of reactions. **A 24 hour grounding period is recommended after each dose providing that aviators are formally briefed about possible delayed reactions.** Individuals who have a past history of urticaria or hypersensitivity phenomena should remain under the previous guidelines (3-5-3 grounding).

JE-Vax is no longer produced. The DoD stockpile is projected to be exhausted in April 2010 and has an expiration date of May 2011. Ixiaro is an available alternative to JE-Vax.

Ixiaro. Ixiaro is administered as a 0.5 mL IM dose. Two doses are given 28 days apart (days 0 and 28). The protective antibody response is 95% at six months and 83.4% at 12 months. Ixiaro is associated with a moderate frequency of mild systemic and local side effects. About 20% of recipients experience headache, 15% experience myalgia, and 50% experience mild local reactions in rates comparable to placebo. **A 12 hour grounding period is recommended for this vaccination.**

Summary- Either JE-Vax or Ixiaro can be used for aviators as described above. JE-Vax will become unavailable by May 2011 unless a shelf-life extension is approved. Ixiaro is likely to have less serious adverse events than JE-Vax. Ixiaro requires only two doses, and requires a 12-hour grounding period instead of 24-hours for JE-Vax. Ixiaro is the preferred vaccine for aviation, especially when the duration of the grounding period impacts mission accomplishment.

MEASLES/MUMPS/RUBELLA (MMR)

This vaccine, composed of live, attenuated viruses, is indicated in adults born after 1956 without a history of documented measles or measles/mumps vaccination. Some people vaccinated before 1980, especially if before 14 months of age, may be inadequately protected and now require revaccination. Contraindications include pregnancy, immunosuppression (except HIV), recent IG administration, or anaphylactic reactions to the immunization, eggs, or neomycin. Efficacy is 95% for all three components. Serious adverse events are rare, but include acute encephalopathy, parotiditis, and orchitis. Transient arthralgias may occur in up to half of first-time recipients, but arthritis and arthropathy are rare. About 5-15% of vaccine recipients have fever up to 21 days post-vaccination and 5% may develop a rash. One study assessed the incidence of adverse events after revaccination. This study noted local injection site discomfort

and flu-like symptoms amongst 6.6% and 3.4% of male and female students respectively. The 4% rate of joint related complaints after revaccination was less than that found after primary vaccination. A 12 hour grounding period is recommended for this vaccination.

MENINGOCOCCAL

Each year, approximately 2,600 people contract meningococcal disease. Of these, 10 to 15% die. Of those who live, another 11 to 19% lose their arms or legs, become deaf, have problems with their nervous system, become mentally retarded, or suffer from seizures or strokes. The meningococcal vaccine is a polysaccharide vaccine that can prevent 4 types of meningococcal disease including 2 of the 3 most common in the United States and a type that causes epidemics in Africa. It is administered as a 0.5 cc SC dose, and is recommended for all children at their preadolescent visit, military recruits, college freshman living in dormitories, microbiologists who might be exposed to the bacteria, anyone with an immune system disorder, asplenic patients, people who might have been exposed to meningitis during an outbreak, and anyone traveling to or living in a part of the world where meningococcal disease is common. Approximately half of vaccine recipients experience mild side effects, such as pain or redness at the injection site. A small percentage of patients also develop fever. Although rare, serious allergic reactions can develop within a few minutes to hours of vaccination. Of note, a few cases of Guillain-Barre syndrome have been reported among people who received the MCV4 vaccine, however there is currently not enough information to determine if this was caused by the vaccine. A 12 hour grounding period is recommended for this vaccination.

PLAGUE

This vaccine is composed of a suspension of killed bacteria, and is given as a dose of 1.0 cc IM. It is used in laboratory workers and travelers to endemic areas. The vaccine is given as a series with a primary dose as above, then 0.2 cc IM doses at 4 weeks and 6 months. Boosters are given every 6 to 12 months as long as exposure continues. There is a 90 to 93% antibody response however efficacy is uncertain. Up to 10% of recipients will develop local reactions. Sterile abscesses and hypersensitivities have also been reported.

PNEUMOVAX (PPV23)

This vaccine was designed to decrease the risk of pneumococcal infection in susceptible individuals such as military recruits, asplenic patients, immunosuppressed individuals, and those over 65. This preparation consists of purified polysaccharide coats of 23 serotypes and is considered to be 60 to 80% efficacious, reducing serious sequelae of infection by about 50%. In asplenic patients it is about 13 -33% effective in producing a two-fold increase in antibody titer. The dose is 0.5 cc IM or SC, and a booster is recommended in high-risk (transplant, nephrotic syndrome, asplenic) individuals at 6 years. Pneumovax has been associated with a 50% local reaction rate, an arthus-like reaction with booster doses, and rarely, anaphylaxis. A 12 hour grounding period is recommended for this vaccination.

POLIO

The inactivated polio virus (IPV) is given as a dose of 0.5 cc IM or SC. The use of oral polio vaccine (OPV) is no longer recommended. Travelers to endemic areas who have received primary immunization during childhood should consider a single booster (IPV) in adulthood, while those who were never vaccinated should be vaccinated according to current CDC guidelines. A 12 hour grounding period is recommended for this vaccination.

SMALLPOX

BACKGROUND: The World Health Organization effectively used smallpox vaccine to eradicate natural smallpox from the planet however regimes hostile to the United States may possess strains of the smallpox virus for use as a biological weapon. While routine vaccination is not recommended for the general population, military and other personnel who serve in high risk parts of the world may receive smallpox vaccine to protect them from the disease in the event of a biological attack.

Expect more side effects within the vaccinated population than normally seen with other vaccines. One expert stated that approximately 10% of vaccine recipients may have side effects significant enough to cause possible distraction during flying activities. The time range for development of side effects varies from day 0 until day 14, with most occurring within 3 to 7 days post-vaccination.

Grounding Period: In view of the complications seen with the smallpox vaccination, a 24 hour grounding period is required. It is recognized that complications from the immunization are most likely in the 3 to 7 day period post immunization. For this reason, close observation and follow-up is recommended by the Designated Aeromedical personnel or health care provider. Personnel should be specifically briefed to report any symptoms or complications during this 3 to 7 day period and to have them evaluated. Depending on the severity, the Flight Surgeon (or Aeromedical Examiner or Aeromedical Physician Assistant) may ground the aviator until symptoms have resolved.

ADDITIONAL INFORMATION: Please review the attached "[Smallpox Fact Sheet - Information for Clinicians](#)" and visit the [CDC web site](#) and military smallpox website (<http://www.vaccines.mil/>) for additional information. Use the [CDC Smallpox Adverse Event Reporting](#) web site to report any adverse events resulting from the administration of the smallpox vaccination.

TYPHOID AND ORAL TYPHOID:

Vaccine is made of a killed suspension of the bacteria, or a new oral 4 dose preparation. The injection is a 0.5 cc IM dose at zero and four weeks with about 50-76% efficacy, and protection for travelers to endemic areas lasts only a few months. This is contrasted with the oral form, which is equally efficacious but confers immunity to the 21a strain that lasts for years (booster required at least every 4 years). It is given every other day before meals for a total of 4 doses, and must be kept refrigerated. Errors in compliance reached 30% of individuals in one study, so direct observation may be the way to go. Adverse reactions to typhoid injections include frequent fever, local swelling and pain, and consequently require a 12 hour downing period. There are no reactions reported to the oral typhoid, therefore no grounding is necessary.

YELLOW FEVER:

This vaccine is used to prevent infection with this flavivirus and its subsequent jaundice, hemorrhage, and albuminuria in travelers to endemic areas (e.g. South America and Africa). It is given as a 0.5 cc SC dose. Booster vaccinations are recommended every 10 years. Efficacy is noted to be high, but adverse side effects include encephalitis/encephalopathy (though fewer than one in a million cases), and anaphylaxis in those individuals allergic to eggs. A 12 hour grounding period is recommended for this vaccination.

COMBINED ADMINISTRATION OF VACCINATIONS:

A number of these vaccines can be given together. Generally, any live virus vaccine can be given with any killed agent or toxoid as long as they are given at the same time and in different anatomic locations. For example, typhoid may be given with either plague or yellow fever. Hepatitis A and yellow fever may be given in the same session. One exception to this is cholera and yellow fever. Administration of these vaccines within 3 weeks of one another results in a poor antibody response. Unless there is insufficient time, 3 to 4 weeks between live virus vaccinations should be sought for maximal antibody production. If possible, vaccines frequently associated with systemic side effects (cholera, typhoid and plague) should not be given simultaneously so that toxicities will not overlap and that a causative agent can be determined should a reaction occur.

PREGNANCY AND VACCINATIONS:

Refer to specific immunization guidelines for vaccination recommendations and precautions during pregnancy.

GROUNDING FOR VACCINATIONS:

OPNAVINST 3710.7 series requires a 12 hour grounding period following immunizations unless otherwise specified in this document. The specific guidelines and grounding periods for each vaccination are described above. As per MANMED Article 15-77, the administration of routine immunizations that require a temporary grounding, do not require issuance of an Aeromedical Grounding Notice. This is a "self-limited" grounding period allowed in the absence of adverse side effects.

18.6 MISCELLANEOUS MEDICATIONS

Last Reviewed: Dec 15

Last Revised: Oct 15

ALLOPURINOL: CD. Waivers are recommended to SG3, Class II, or Class III. Re-evaluation for upgrade from SG3 to SG1 is considered in 3 months if member remains asymptomatic and on a stable dose of medication.

ANTI-HISTAMINES (SEDATING):

CD. The member should be grounded for the duration of therapy.

ANTI-HISTAMINES (NON-SEDATING):

NCD. [Allegra](#) and [Claritin](#) are NCD if given in accordance with the [Allergic/Vasomotor Rhinitis](#) section of the Waiver Guide. Refer to this section for additional restrictions and clarification. [Zyrtec](#), although considered by some to be non-sedating, still has a moderate sedating effect and is therefore not approved (CD) for use in aviation personnel.

CLOMIPHENE (CLOMID): CD- No Waivers.

CONTRACEPTIVES:

DEPO-PROVERA : NCD. Any grounding period at discretion of the local Flight Surgeon to assure tolerance.

LEVONORGESTEROL (NORPLANT): NCD. Any grounding period at discretion of the local Flight Surgeon to assure tolerance.

PROGESTASERT IUD: NCD. Any grounding period at discretion of the local Flight Surgeon to assure tolerance.

DECONGESTANTS: CD. Requires temporary grounding while in use.

FINASTERIDE (PROPECIA/PROSCAR):

CD. A waiver can be considered after a two week grounding. If the patient remains asymptomatic, a LBFS may issue an up chit. [Finasteride](#) may be utilized for prostatic hypertrophy or alopecia. DoD pharmacy does not allow prescriptions of [finasteride](#) for hair loss.

H2 BLOCKERS:

RANITIDINE, CIMETIDINE, FAMOTIDINE: CD. A waiver is required for any chronic use. Refer to the Gastroenterology Waiver Guide section on [reflux esophagitis](#) for additional information.

INHALED STEROIDS: CD. Decisions are individualized. Any chronic use requires a waiver. Call NAMI Code 53HN for additional guidance.

ISOTRETINOIN (ACUTANE, AMNESTEEM, CLARAVIS, SOTRET): CD. No waiver.

Resumption of flight status is permitted after member is off medication for 3 months, has a normal slit lamp exam, and triglyceride levels are documented as normal. Cystic acne, if severe enough to need [Accutane](#), may be disqualifying. The 3 month delay after

cessation of treatment also allows for an evaluation of how the member does without the medication.

LEVOTHYROXINE (SYNTHROID): CD. A waiver may be requested when member is clinically and chemically euthyroid on stable dosage.

LINDANE (K WELL): NCD. Requires a 48-hour grounding period. Kwell can be absorbed in variable amounts and give some significant CNS side effects. Aviation personnel must be grounded for 48 hours after the compound is washed off.

MESALAMINE (ASACOL, ROWASA, ETC.): CD. A major advantage of mesalamine is that it avoids some side effects associated with the sulfapyridine moiety of sulfasalazine. Waiver will be considered after maintaining clinical remission for one month without evidence of side effects.

MINOXIDIL (TOPICAL): CD. No Waiver.

NEDOCROMIL (TILADE): CD. Tilade may be considered for waiver for in designated aviation personnel for the preventive treatment of mild to moderate asthma or cold-induced and exercise-induced bronchospasm. Member will be eligible for waiver consideration and return to flight status at a minimum two weeks after remaining symptom free on a stable dose of medication with demonstrated normal pulmonary function tests. Waivers are restricted to non-high performance aircraft.

NASAL STEROIDS:

FLONASE, NASONEX, RHINOCORT: NCD. Refer to the [Allergic/Vasomotor Rhinitis](#) section of the Waiver Guide under Ear/Nose/Throat for additional restrictions and clarification.

SMOKING CESSATION:

NICORETTE GUM: NCD if the following conditions are met:

1. Enrolled in formal organized stop smoking program.
2. Close observation by flight surgeon.
3. No adverse effects.
4. Duration of use does not exceed three months.

NICOTINE TRANSDERMAL SYSTEM (NICODERM): NCD. Aviators should be grounded for 48 hours following application of first patch.

All other medications for tobacco cessation are not approved for use by personnel on active flight status, so require grounding during treatment. This can often be planned to coincide with non-flying periods. Guidance for timing of return to flight is based on the elimination half-life of the drug being used, as follows:

VARENICLINE (CHANTIX®): Varenicline has an elimination half-life of 24 hours, so individuals should be grounded for one more week after finishing Chantix

BUPROPION (ZYBAN®, WELLBUTRIN®): Bupropion is cleared more quickly, but only about 1% is excreted unchanged in the urine; the rest is metabolized to three major active

metabolites, threohydrobupropion, erythrohydrobupropion and hydroxybupropion, which accumulate to levels much higher than the parent compound and can have extended half-lives of as long as 43 hours. Individuals taking bupropion should therefore be kept down for two weeks following completion of treatment.

NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDS):

ASPIRIN: NCD for occasional analgesic use or at cardioprotective dosing. Other chronic use is CD and requires a waiver.

IBUPROFEN/NAPROXEN: NCD. Medication can be used for short term use under direct supervision of Flight Surgeon. Any chronic or high dose use is disqualifying. If recommending that an aviator continue to fly during treatment, consider the underlying reason for its use. It may be the condition which is disqualifying.

INDOCIN: CD. No waiver. Ground during medication use and for two weeks after medication is completed.

PHOSPHODIESTERASE INHIBITORS (PDI):

SILDENAFIL (VIAGRA), VARDENAFIL (LEVITRA), TADALAFIL (CIALIS):

1. Long-acting PDI, tadalafil (Cialis): CD. No Waivers
2. Short-acting PDI: sildenafil (Viagra), vardenafil (Levitra): CD. Waivers possible for:
 - a. Initial Applicants – considered on a case-by-case basis
 - b. Designated Personnel – may request waiver after evaluation for the cause of ED.
3. Information Required:
 - a. AMS – history, physical, lab, prior treatment course, side effect.
 - b. Consultation – Urology, Internal or Family Medicine – Must evaluate causes of ED including co-morbid conditions such as cardiovascular (hypertension, atherosclerosis, or hyperlipidemia), diabetes mellitus, depression and alcoholism
 - c. An exercise stress test should be completed prior to waiver submission whenever indicated. If test is not performed, reasons should be substantiated in AMS or by consultants.
4. Follow-up: annual – to assess efficacy, side effects, and significant changes in health status including medications
5. Treatment:
 - a. Must be free of side-effect for 2 doses after beginning medication before returning to flying duty
 - b. No flying duties within 12 hours of last dose (medicine use to briefing time)
6. Links:
 - a. Diagnostic evaluation: [Link to diagnostic evaluation](http://www.aafp.org/afp/2000/0101/p95.html) (<http://www.aafp.org/afp/2000/0101/p95.html>)
 - b. Precautions: [Link to precautions](http://www.aafp.org/afp/1999/0915/p1159.html) (<http://www.aafp.org/afp/1999/0915/p1159.html>)
 - c. Medications: sildenafil ([Viagra®](#)), vardenafil ([Levitra®](#))

PROBENECID:

CD. Waiver is required for any long term treatment.

PROTON PUMP INHIBITORS:

OMEPRAZOLE, LANSOPRAZOLE, RABEPRAZOLE: CD. Waiver required for chronic use. Refer to the Gastroenterology Waiver Guide section on [reflux esophagitis](#) for additional information.

SUCRALFATE (CARAFATE):

NCD when used in dosages of 1 gm bid or less. However, the diagnosis of peptic ulcer disease is certainly CD and requires a waiver.

SULFASALAZINE (AZULFIDINE):

CD. Waiver considered after maintaining clinical remission for one month without evidence of side effects.

TAMOXIFEN: CD. No Waiver.

TOPICAL COMPOUNDS:

As a general rule, medications applied to the surface of the body which are not absorbed to any significant extent are NCD. However, please see notes on [Kwell](#). The recommended initial treatment is over a weekend to allow return to flight duties the following Monday, thus minimizing flight schedule loss.